



Delta activity independent of its activity as a ligand of Notch.

Lee-Peng Mok, Tielin Qin, Boris Bardot, Matthew Lecomte, Asal Homayouni, Francois Ahimou, Cedric Wesley

► To cite this version:

Lee-Peng Mok, Tielin Qin, Boris Bardot, Matthew Lecomte, Asal Homayouni, et al.. Delta activity independent of its activity as a ligand of Notch.. BMC Developmental Biology, 2005, 5, pp.6. 10.1186/1471-213X-5-6 . inserm-00092484

HAL Id: inserm-00092484

<https://www.hal.inserm.fr/inserm-00092484>

Submitted on 12 Sep 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research article

Open Access

Delta activity independent of its activity as a ligand of Notch

Lee-Peng Mok¹, Tielin Qin², Boris Bardot³, Matthew LeComte¹,
Asal Homayouni⁴, Francois Ahimou⁵ and Cedric Wesley^{*1}

Address: ¹Department of Microbiology and Molecular Genetics, 322 Stafford Hall, 95 Carrigan Drive, The University of Vermont, Burlington, VT 05405, USA, ²Rollins School of Public Health, Emory University, Atlanta, GA 30322, USA, ³INSERM E365, Faculte de Medecine Lariboisiere, 10 avenue de Verdun, Paris 75010, France, ⁴College of Veterinary Medicine, Kansas State University, Manhattan, KS 66506, USA and ⁵Department of Civil Engineering, The University of Minnesota, Minneapolis, MN 55455, USA

Email: Lee-Peng Mok - lmok@uvm.edu; Tielin Qin - tqin@sph.emory.edu; Boris Bardot - bbardot@uvm.edu; Matthew LeComte - Matthew.LeComte@uvm.edu; Asal Homayouni - ahomayou@vet.k-state.edu; Francois Ahimou - ahimo001@tc.umn.edu; Cedric Wesley* - cwesley@uvm.edu

* Corresponding author

Published: 10 March 2005

Received: 15 August 2004

BMC Developmental Biology 2005, 5:6 doi:10.1186/1471-213X-5-6

Accepted: 10 March 2005

This article is available from: <http://www.biomedcentral.com/1471-213X/5/6>

© 2005 Mok et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Delta, Notch, and Scabrous often function together to make different cell types and refine tissue patterns during *Drosophila* development. Delta is known as the ligand that triggers Notch receptor activity. Scabrous is known to bind Notch and promote Notch activity in response to Delta. It is not known if Scabrous binds Delta or Delta has activity other than its activity as a ligand of Notch. It is very difficult to clearly determine this binding or activity *in vivo* as all Notch, Delta, and Scabrous activities are required simultaneously or successively in an inter-dependent manner.

Results: Using *Drosophila* cultured cells we show that the full length Delta promotes accumulation of Daughterless protein, *fringe* RNA, and *pangolin* RNA in the absence of Scabrous or Notch. Scabrous binds Delta and suppresses this activity even though it increases the level of the Delta intracellular domain. We also show that Scabrous can promote Notch receptor activity, in the absence of Delta.

Conclusion: Delta has activity that is independent of its activity as a ligand of Notch. Scabrous suppresses this Delta activity. Scabrous also promotes Notch activity that is dependent on Delta's ligand activity. Thus, Notch, Delta, and Scabrous might function in complex combinatorial or mutually exclusive interactions during development. The data reported here will be of significant help in understanding these interactions *in vivo*.

Background

Notch (N) and Delta (DI) are cell surface proteins that are required for differentiation of almost all tissues in the fruit fly *Drosophila melanogaster*. They are evolutionarily conserved, functioning similarly in animals from worms to humans [1,2]. The best-known instance of their function is the process of lateral inhibition that initiates differenti-

ation of the neuronal and epidermal tissues from proneural cells that are predisposed to making the neuronal tissue. Proneural cells express high levels of the neuronal transcription co-factors from the Achaete Scute Complex (ASC) or related genes [3,4]. These factors require their partner Daughterless (Da) to activate transcription of the neurogenesis genes [5-7]. Da is expressed at low levels in

all *Drosophila* cells [8] and up regulated in proneural cells specified to differentiate the neurons [5]. Whether or not the up regulation of Da expression is part of lateral inhibition is not clear in *Drosophila*. In *Caenorhabditis elegans*, however, the differential accumulation of the Da homolog HLH-2 is the earliest detectable difference between the cells taking up alternate fates during lateral inhibition [9]. As N and Dl are known to regulate Da expression [10], it is very possible that Da expression is regulated during lateral inhibition in flies as well.

When N expressed on one proneural cell binds Dl expressed on the neighboring proneural cell, N is proteolytically cleaved to release the Notch intracellular domain (N^{intra}) from the plasma membrane. N^{intra} translocates to the nucleus and, in association with the transcription factor Suppressor of Hairless (SuH), activates transcription of the *Enhancer of split Complex* (*E(spl)C*) genes. Cells that express a high level of *E(spl)C* RNA suppress their neuronal predisposition, become the epidermal precursor cells (EPCs), and differentiate the epidermis. Cells that express a low level of *E(spl)C* RNA and a high level of Da protein become the Neuronal Precursor Cells (NPCs) and differentiate the nervous system [1,2,5,11]. From here onwards, we refer to this SuH dependent N activity that promotes expression of *E(spl)C* RNA as SuH/N^{intra} signaling. A 1.5 to 2-fold difference in the level of SuH/N^{intra} signaling is sufficient to initiate specification of the EPCs and the NPCs [11]. This difference is amplified by subsequent activities of N and Dl, or activities of other genes responding to the initial difference in the level of SuH/N^{intra} signaling. The lateral inhibition process described above is repeatedly used during development for differentiation of various tissues with minor variations or changes in target genes.

Scabrous (Sca) is a secreted factor that is produced at high levels in the NPCs and functions non-autonomously to promote specification of the EPCs during differentiation of the compound eye and the bristle organ [12,13]. In its absence, lateral inhibition is not abolished but is reduced in strength or becomes imprecise indicating that Sca only refines the process. Sca binds N and stabilizes it. These actions promote formation of sharp boundaries between neuronal and non-neuronal cells during development of the compound eye [14]. The possibility that Sca might bind Dl as well is suggested by the observation that simultaneous over expression of Sca almost completely blocks the effect of Dl over-expression on wing margin development but hardly modifies the effect of N over-expression [15]. Dl and Sca have also been observed to co-localize in intracellular vesicles *in vivo* [13]. The observations that Sca can promote N activity [14] but block Dl activity are paradoxical as SuH/N^{intra} signaling is very much dependent on the activities of both N and Dl. One explanation for this paradox could be, that Sca promotes lateral inhibition

by having one effect through N and a different one through Dl. Therefore, we addressed the following questions in this study. Does Sca bind Dl? If yes, does it affect any Dl activity? Are there Dl activities independent of its activity as a ligand of N? Is Sca capable of activating N in the absence of Dl?


N and Dl are expressed in almost all cells *in vivo* and N receptor activities in response to Dl binding are widely used during development. In developmental instances where Sca is present, the expression data suggest that both N and Dl will have access to Sca. Thus, it is very difficult to separate *in vivo* the activities of N alone, Dl alone, N on Dl, Dl on N, Sca on N, Sca on Dl, and Sca on N and Dl together. Therefore, we addressed the questions posed above in an *in vitro* model system based on *Drosophila* Schneider (S2) cells. S2 cells do not express the endogenous N, Dl, or Sca [14,16]. S2 cells expressing N (S2-N cells) mixed with S2 cells expressing Dl (S2-Dl cells) reproduce all aspects of lateral inhibition [16-22]. Using these cells and the medium prepared from S2 cells expressing Sca into the medium [14], we show that Sca binds Dl, Dl has activity independent of its activity as a ligand of N, Sca can affect this activity of Dl, and Sca can activate N in the absence of Dl. These observations would be useful for undertaking the challenging task of determining how the various activities of N, Dl, and Sca are integrated during tissue differentiation.

Results

Sca associates with Dl

Although N and Sca complexes could be immuno-precipitated [14], we, and others [23], had failed to detect Sca on S2-N cells. We suspected that some factor present in the tissue culture medium was washed away when the cells were processed for immuno-fluorescent detection of Sca. To overcome such problems, we made Sca-GFP and established stable S2 cells expressing it (S2-Sca-GFP cells). S2-Sca-GFP cells produced the Sca-GFP protein of the expected size (as determined by western blotting) and both Sca and GFP antibodies recognized this protein (data not shown). We concluded that S2-Sca-GFP cells expressed the expected Sca-GFP protein and used the conditioned medium from these cells to treat live S2-N, S2-Dl, and S2 cells.

Live S2-Dl cells showed the strongest GFP signals, followed by live S2-N cells, and then live S2 cells (Fig. 1A-C). The signals were so strong on the S2-Dl cells that the signals on S2-N cells were not obvious at the same brightness/contrast settings. When cells were simultaneously fixed and rinsed with 1X PBS, the signals were comparable at the same settings (insets in Fig. 1A-C). Signals could not be detected on S2-N or S2-Dl cells after three 5-minute washes with 1X PBS, confirming our suspicion that the

**Figure 1**

Sca associates with DI. **A-C.** Fluorescent photomicrographs of different cell lines treated with Sca-GFP medium for 30 minutes. Cells simultaneously fixed and rinsed in 4% paraformaldehyde/1X PBS are shown in the insets. Experiments were repeated three times. **D.** Western blots showing recovery of DI in Sca immuno-precipitates from total protein extracts prepared from S2-DI cells treated with S2-Sca cells. S2-Sca cells were used instead of Sca conditioned medium to maximize the ratio of bound to unbound Sca. Cross-linker = membrane insoluble and cleavable 3,3'- Dithiobis (sulfosuccinimidylpropionate) (DTSSP), which cross-links proteins interacting at the cell surface. IP Ab = antibody used for immunoprecipitation; W Ab = antibody used on the western blot; ppt = immunoprecipitate; super = supernatant. Experiments were repeated two times.

standard immuno-fluorescence procedure is inappropriate for detecting Sca binding on S2-N or S2-Dl cells. Secreted GFP did not bind the surfaces of any of these cells (data not shown). This indicated that the Sca part of Sca-GFP fusion protein bound the S2-N and S2-Dl cell surfaces. In all experiments conducted to determine the activity of Sca, N, or Dl, that are described below, we used only S2 cells expressing the wild type Sca because (1) we do not perform washes to remove non-specifically bound proteins and (2) we wanted to avoid possible GFP associated effects (stability, etc.).

We have previously shown that Sca and N form complexes [14]. To determine whether Sca forms complexes with Dl, we performed immuno-precipitation experiments with S2-Dl cells that were co-cultured with S2-Sca cells. Proteins interacting at the cell surfaces were either cross-linked or un-linked prior to cell lysis for protein extraction. Membrane insoluble cross-linkers improve recovery of cell surface complexes [18,24]. Sca immuno-precipitation recovered Dl strongly in the presence of cross-linkers and relatively weakly in the absence of cross-linkers (Fig. 1D). No bands were observed when S2 cells were used instead of S2-Dl cells (data not shown). In the reverse experiments, Dl immuno-precipitations failed to recover Sca, possibly because there was too much unbound Dl in the extracts. Dl and Sca were not detected in the absence of immuno-precipitation antibodies (Fig. 1D, lanes 1 and 4) or in the absence of Scabrous (data not shown). We also recovered Sca in Dl immuno-precipitations and Dl in Sca immuno-precipitations from protein extracts of wildtype embryos (data not shown). These observations indicated that Sca associates with Dl. We explored the consequence of this association.

Da expression in Dl cells is reduced in response to Sca

N promotes expression of *E(spl)C m3* gene in response to Dl [19,20]. We examined whether Sca promoted expression of *E(spl)C m3* in S-N cells or S2-Dl cells and found that it was indeed the case with S2-N cells, but not with S2-Dl cells (Fig. 2A, lanes 1–6). S2-N cells showed a low level of *E(spl)C m3* expression when S2-Dl or S2-DlΔI cells were replaced with S2 cells, in the absence of Sca (Fig. 2A, lanes 1, 7–8); S2-Dl or S2-DlΔI cells mixed with S2 cells did not show any accumulation (Fig. 2A, lanes 13–16). The low level of *E(spl)C m3* RNA expression in S2-N cells in the absence of ligands is due to the low level of N^{intra} produced upon induction of N expression in S2 cells [18]. This expression increases upon ligand treatment [18], resulting in increased expression of *E(spl)C m3* RNA expression (Fig. 2A, lanes 2, 10, 12). Numerous repetitions of the experiments indicated that Dl is a more potent ligand of N than Sca with respect to induction of *E(spl)C m3* expression (data not shown).

E(spl)C m3 expression appeared to be solely dependent on N activation and the Notch intracellular domain as it was promoted in S2-N cells treated with either S2-Dl cells or S2-DlΔI cells (Fig. 2A, lanes 7–12). As DlΔI lacks the intracellular domain, it is expected to behave only as a ligand of N and not generate any intracellular signal of its own in response to N binding. We observe comparable levels of SuH/N^{intra} signaling with S2-Dl and S2-DlΔI cells (Fig. 2A, lanes 10, 12). This is not consistent with the *in vivo* findings that the Dl intracellular domain (lacking in DlΔI) is required for SuH/N^{intra} signaling, possibly for promoting Dl internalization that results in exerting a 'pull' on N and increased production of N^{intra} [21,25–29]. However, our results are consistent with other S2 cell studies showing that even fixed S2-Dl cells can promote production of SuH/N^{intra} signaling in S2-N cells [19]. Thus, it is possible that that Dl internalization and pulling is not required for SuH/N^{intra} signaling in S2 cells. In any case, in our S2 cell system, the S2-N and S2-Dl cells require shaking for formation of cell aggregates. As a consequence, we shake all cell mixtures, including those containing the secreted ligand and Sca. This shaking might have simulated the pulling effect and overcome any deficiency DlΔI might have in this regard thereby resulting in a level of SuH/N^{intra} signaling that is comparable to that produced by the full length Dl.

We examined the expression of various proteins known to be involved in lateral inhibition to find out if Dl expression affected them. They were Numb, Dishevelled, Suppressor of Hairless, Wingless, Hairless, Hairless, Achaete, Da, and Armadillo. We found a relatively high level of Da protein in S2-Dl cells compared with the level in S2-DlΔI cells (Fig. 2B, lanes 1–2). Similar levels of Da were expressed in S2-DlΔI and S2 cells (data not shown). Two independently transfected S2-Dl cell lines also showed high levels of Da, and un-induced S2-Dl cells showed background levels of Da, indicating that Dl expression promotes Da expression (Fig. 2B, lanes 3–6). Increase in Da levels appeared to be specifically linked to Dl expression, as S2-N cells did not show an increase (Fig. 2C, compare lanes 1 & 3). Overall, Da expression in S2-Dl cells was 2.18X higher (+/- 0.37, *p* < 0.05) than the level in S2 cells, sometimes more than 5X higher. Here and in all cases to follow, the blots shown in the figures are the most representative blots among replications. Graphs show quantification, relative to standards or other proteins (as indicated), of signals on the blots composing the figures as the response can be assessed only in comparison to the control lanes in the same experiment. Pooling data from all replications of an experiment obscured the response, or misrepresented the data, due to variation between different batches of cells. Therefore, we computed error variance for the degree of response over all replications of an experiment. These values for important responses are mentioned in the text.




Figure 2
DI down-regulates Daughterless protein expression, and N up-regulates *E(spl)C m3* gene expression, in response to Scabrous. **A.** Northern blots of total RNA from the indicated cell mixtures extracted at 0 or 45 minutes after treatment with medium containing Sca (+) or not (-). Gene probes used are shown on the right. m3 = *E(spl)C m3* and rp 49 = a ribosomal protein gene used to show the levels of total RNA in the lanes in all northern blots. Sca = conditioned medium prepared from the S2-Sca stable cell line in all experiments here onwards. The control medium used along side Sca medium (-) was prepared from heat shocked S2 cells. Experiments were repeated two times. For unknown reasons, the medium collected from heat shocked S2 cells (used in lanes 1, 3, and 5) produced higher background levels of *E(spl)C m3* RNA in S2-N cells (lane 1). **B.** Western blots showing the levels of Da and DI in different DI cell lines. S2-DI, S2-DI(1) and S2-DI(2) are independently established hsDI cell lines. Ui = un-induced (i.e., not heat shocked). Hsp70 = heat shock 70 protein used to show the levels of proteins in the lanes of all western blots. DI and DIΔI were detected with αDIEC. Da signals here (and the indicated signals elsewhere) were quantified relative to Hsp70 (western blots), rp49 (northern blots), or other indicated molecules, using the NIH Image 1.63 program. These experiments were repeated more than ten times. **C.** Western blots showing Da levels in the indicated cell mixtures, with (+) or without (-) Sca. These experiments were repeated five times.

The number of repetitions of an experiment is indicated in the figure legends.

We examined Da levels in S2-Dl and S2-N cells that were treated or not treated with Sca conditioned medium. We found that Sca treatment decreased the levels of Da in S2-Dl cells (Fig. 2C, lanes 3–4). The levels in S2-N cells were low and unaffected by Sca treatment (Fig. 2C, lanes 1–2). These experiments suggested that Sca blocks accumulation of Da in S2-Dl cells (2.81X, \pm 0.59, $p < 0.05$). We also determined the levels of Da when S2-N and S2-Dl cells were together in the absence of Sca. The level of Da never increased (Fig. 2C, lanes 6–7). As N activation suppresses *daughterless* RNA expression [18], it was possible that N activation suppressed Da expression in S2-N cells and masked an increase in S2-Dl cells. To determine if this was the case, we compared the level of Da in mixtures of S2-DlΔI cells and S2 cells with mixtures of S2-DlΔI cells and S2-N cells. As S2-Dl and S2-DlΔI cells activate N equally well (see Fig. 2A, lanes 9–12), any change in Da level would be due to N activation. We found comparable levels of Da in the two samples (Fig. 2C, lanes 8–9). Thus, S2-Dl cells do not appear to increase Da expression in response to S2-N cells.

When S2-N and S2-Dl cells were together in the presence of Sca, the levels of Da protein and *E(spl)C m3* RNA were very variable (data not shown). This was possibly due to the varying combinations of Sca effect on S2-Dl cells, Sca effect on S2-N cells, Dl effect on S2-N cells, and N effect on S2-Dl cells.

Dl is processed to produce Dl intracellular domain (DIIC), constitutively, and the levels of DIIC increase upon N treatment [30–33]. Therefore, we examined the levels of DIIC following treatment of S2-Dl cells with S2-N cells or Sca medium. We found that the DIIC levels increased by 25 to 50% (relative to Dl levels) with both treatments (Fig. 3A and 3B). We examined the levels of Da in S2 cells expressing DIIC and DIIMIC (lacking the extracellular domain only and including the transmembrane domain). The levels of Da in S2-DIIC and S2-DIIMIC were always comparable to or lower than the level in S2 or S2-DlΔI cells (Fig. 3C). Also, we found Da levels to be negatively correlated with the accumulation of DIIC in time-course experiments (Fig. 3D). This negative correlation could be a direct consequence of the accumulation of DIIC or due to autoregulation of the *da* gene [34]. We examined the levels of Da in flies expressing heat shock induced Dl, N, or Sca, in flies heterozygous for the null alleles of N or Dl, and in flies homozygous for a null allele of Sca. We found that Da expression was strongly associated with Dl expression rather than with N expression, and inconsistently associated with Sca expression. These data are consistent with our findings in S2 cells but are not

shown, as we cannot clearly separate the effects of N, Dl, and Sca, the way we can do in S2 cells. The experiments described in this section indicate that Da accumulation is promoted by the full-length Dl, not by the Dl intracellular domains (DIIC or DIIMIC), and Sca suppresses the activity of the full length Dl. The experiments also indicate that Sca promotes *E(spl)C m3* RNA expression in S2-N cells even in the absence of Dl.

Dl regulates expression of fringe and pangolin

To gather additional evidence for Dl activity independent of its activity as a ligand of Notch, we performed microarray experiments using the Affymetrix Drosophila GeneChip Arrays to compare gene expression in S2 cells and S2-Dl cells. Many genes relevant to known Dl functions responded in S2-Dl cells (at $p < 0.05$, $n = 3 \times 2$ pooled samples): axonal path finding genes (e.g., *Gef64C*, 39.38XUp; *Tenascin major*, 6.77XUp), actin-based cell motility and kinases (*Rho-Kinase*, 15.08XUp; *Rhopilin* 3.4XUp; *nemo* 1.72XUp, *basket* 1.69XUp; *pointed* 2.2XUp), N signaling pathway genes (e.g., *reaper*, 2.26XUp; *sanpodo*, 1.91XUp), and oogenesis genes (e.g., *swallow*, 8.12XUp; *sprouty*, 3.58XUp). Expression of *transformer*, was also up (1.76X) and it is significant in the light of our observation that Dl promotes expression of Da: both Da and *transformer* are involved in sex determination. Expression of *da* RNA was not significantly increased in S2-Dl cells, possibly due to the negative part of the *da* gene autoregulation system [34]. The detailed analyses with validations will be published elsewhere. The experiment also identified *fringe* (*fng*) and *pangolin* (*pan*) as responding to Dl expression. *fng* is a glycosyl transferase that regulates the affinity of N for Dl [35–37], and possibly also the affinity of Dl for N [38]. *pan* is a transcription factor functioning in the Wingless (Wg) pathway [39,40]. Notch and Wg pathways interact closely at many differentiation events during development [24,41–44]. Therefore, we chose *fng* and *pan* for further investigation.

Northern blot analyses showed that the expression of *fng* and *pan* was higher in S2-Dl cells compared with S2-N or S2-DlΔI cells (Fig. 4A). DIIC and DIIMIC promoted expression of *pan* and *fng* weakly, if at all (Fig. 4B). Two independently established S2-Dl cell lines also showed higher levels of *fng* and *pan* RNAs (Fig. 4C). Sca treatment S2-Dl cells reduced the levels of *fng* and *pan* RNA (Fig. 4D). This reduction was expected as Sca reduces the levels of the full length Dl (see Fig. 3). Thus, just as it was the case with Da expression, the full length Dl, not any of its parts, strongly promoted *pan* and *fng* expression. We examined the levels of *fng* and *pan* RNA in flies expressing heat shock induced Dl, N, or Sca, in flies heterozygous for the null alleles of N or Dl, and in flies homozygous for a null allele of Sca. We found that *fng* and *pan* RNA expression was strongly associated with Dl expression





Figure 3
The levels of cleaved DI intracellular domain is not associated with high levels of Da. **A.** Western blots (from a 8% SDS-PAGE) showing the level of DI and DIIC in the indicated cell mixtures, with (+) or without (-) Sca. **B.** Western blots (from a 12% gel) showing the levels of DI and DIIC in S2-DI cells treated medium containing different levels of Sca. **C.** Western blots showing the levels of Da in the indicated cell mixtures. **D.** Western blots showing the levels of Da, DI, and DIIC at different times following heat shock induction of DI in S2-DI cells. All experiments were repeated at least three times.

**Figure 4**

DI promotes expression of *fng* and *pan*. **A**. Northern blots showing *fng* and *pan* expression in the indicated cell mixtures at 0 and 45 minutes after cell mixing. **B**. Northern blots showing *fng* and *pan* expression in the indicated cell lines two hours after induction of expression. **C**. Northern blots showing *fng* and *pan* expression in two other independently established S2-DI cell lines. Cells used for lanes 1–2 were uninduced (ui); cells used for lanes 3–4 were heat shock induced. **D**. Northern blots showing *fng* and *pan* expression in S2-DI cells that were either untreated or treated with Sca medium. All experiments were repeated at least three times. The *fng* band marked with an asterisk corresponds to the published mRNA [35]. Only this band was used for *fng* quantification. The *pan* band shown is consistent with the information described in van de Wetering et al. [40] and Brunner et al. [39].

rather than with N expression, and inconsistently associated with Sca expression. These data are consistent with our S2-Dl cells data but are not shown, as we cannot clearly separate the effects of N, Dl, and Sca, the way we can do in S2 cells.

Discussion

Our experiment addressed four questions. Does Sca bind Dl? If yes, does it affect any Dl activity? Are there Dl activities independent of its activity as a ligand of N? Is Sca capable of activating N in the absence of Dl? Results described in Figure 1 show that Sca binds Dl. This binding is not dependent on N as S2-Dl cells do not express N. We have previously shown that Sca binds N [14]. It is possible that Sca binds N or Dl stronger when they are present together on the same cell or on neighboring cells. It would be possible to test this in the future using Atomic Force Microscopy that is best suited for determining binding strengths of cell surface proteins like N or Dl [21]. Results in Figure 2A show that Sca can promote SuH/N^{intra} signaling through N in the absence of Dl, as S2-N cells do not express Dl. However, numerous repetitions of the experiment indicate that Sca is not as potent as Dl in this regard. This is consistent with the fact that lateral inhibition is blocked in the absence of zygotic Dl, which does not affect proneural cluster formation and thereby Sca expression. It would have been relatively easy to determine if overexpression of Sca in the absence of Dl rescues SuH/N^{intra} signaling phenotypes, and the extent of this rescue, if Dl did not have any activity independent of N. Hopefully, it would be possible in the future, when we better understand this Dl activity and are able to circumvent it. Results in Figure 2A also show that the expression of E(spl)C m3 gene, a target of SuH/N^{intra} signaling pathway, is responsive only to N indicating that this pathway is unlikely to be involved in mediating Dl activities.

Results described in Figures 2, 3, 4 and our microarray analysis show that Dl has activity independent of its activity as a ligand of N and Dl could be a receptor of Sca. This is clearly shown in experiments with S2-Dl cells that do not express N and we do not provide either N or Sca (Fig. 2B, C; 4A). The Dl activities we have described- promotion of expression of Da protein, *fng* RNA, and *pan* RNA- can be detected *in vivo* as well although the interpretation here is not simple due to the many possible interactions among N, Dl, and Sca that cannot be easily sorted out. However, these data (which we do not show) strongly suggest that the Dl activities we have described in S2 cells represent the *in vivo* Dl activities during development.

The N independent Dl activity we have described is dependent on the full length Dl, not just on its intracellular domain or the extracellular domain (Figs. 2C; 3; 5A-B). This is different from the situation with N whose activity

is based on the activity of its intracellular domain [45-47]. Accordingly, treatment with Sca, which promotes production of the Dl intracellular domain, suppresses Dl activity related to Da rather than promote it (Figs. 2C; 3). This observation also indicates that Sca is able to affect Dl activities. A clean dissection of Sca effects through N from its effects through Dl would require identification of Sca binding sites on Dl, and N and Dl binding sites on Sca. We know that Dl and Sca bind different regions of N [14,48,49]. It would not be too surprising if Dl bound N and Sca in different regions, and if Sca bound N and D in different regions. With that knowledge and suitable mutants, we might be able to determine whether N, Dl, and Sca activities function in a mutually exclusive or combinatorial manner *in vivo*.

Dl activity that is independent of its N ligand activity has been speculated for some time. Efforts to identify it have intensified since the discovery that Dl gets proteolytically processed in the same manner as N [30-33]. However, it is extremely difficult to separate these two activities of Dl. The proof that the Dl activity we have identified actually functions during development in the expected manner, the details of the mechanisms underlying this function, and a better integration of the known functions of N, Dl, and Sca, will have to await more work which is neither quick nor simple. We hope that this work stimulates more efforts towards this task and makes this task a bit easier by identifying the potential of Sca as a regulator of Dl activity and the possibility that the full length Dl might be important for Dl activity independent of N, or Sca. Sca could also serve as a great tool for *in vivo* dissection of Dl response to N, as Sca and N appear to have a similar effect on Dl. It is quite likely that our experiments did not pick up Dl receptor activity in response to N or Sca. In any case, the potential developmental significance of our findings is briefly discussed below.

Da is a widely expressed protein and cells requiring its function show only a modest increase in its levels [5,6,8] indicating that, just like N^{intra}/SuH signaling, small changes in Da levels might be sufficient for initiating or augmenting NPC specification and promoting neuronal differentiation. Small changes in Da levels might also be imposed by the built-in autoregulation of the *da* gene [34]. According to the well-accepted lateral inhibition model in the field, Dl activity as a ligand of N is postulated to increase in the NPCs and N receptor activity in response to Dl is postulated to increase in the EPCs [11]. Accordingly, Dl expression has been observed to increase in the NPCs or their equivalent cell types in certain instances involving N and Dl functions [50,51]. Our data suggest that an increase in Da levels in these instances could be due to the accumulation of the full length Dl, not any its parts such as DIIC, DII, or DITMIC. The requirement for

the intracellular and the extracellular domains to be linked might mean that we have detected DL activity requiring DL's presence at the membrane or in the cytoplasm. This is consistent with the report that the cellular transformation ability of Jagged 1, a mammalian DL homolog, requires an intact protein containing both the extracellular and the intracellular domains [52]. It is possible that DLIC, in the nucleus [31], promotes other activity that is different from the one described here. It is also possible that *Da*, *fng*, or *pan* might not be the direct target of the full length DL activity. Our microarray data indicate that many other genes (including some in the RAS or EGFR signaling pathways) are strongly up regulated in DL expressing cells. It is possible that one of these genes is the primary target. It is also possible that *Da*, *fng*, or *pan* accumulation is significant only in the context of these other genes. We will have to await validation of other putative targets of DL activity, and evaluation of their role in lateral inhibition or other activities involving N and/or DL, to determine if *Da*, *fng*, or *pan* are typical or atypical targets of DL activity.

N/DL binding and SuH/N^{intra} signaling are strongly affected by the functions of glycosyl transferases such as *fng*. The possibility that DL, and not N, regulates *fng* RNA expression might explain some of the very complex functions of these glycosyl transferases and the complex interactions between N and DL during lateral inhibition. As N and DL activities are known to strongly interact with the Wg signaling pathway, it is interesting that DL promotes *pan* expression. It is possible that DL activity independent of N accounts for some of the interactions between the N and the Wg pathways. So far, these interactions have been considered only from the perspective of N receptor activity.

Lastly, our data suggest interesting interactions among DL, N, and Sca in instances of lateral inhibition and tissue differentiation when their functions overlap. The full length DL promotes *Da* accumulation, not any of its parts that might result from processing in response to N or Sca binding. Thus, both the processed N and DL might promote EPC specification- processed N through *E(spl)C* RNA and processed DL through suppression of *Da* expression. Consequently, lateral inhibition might initiate with symmetrical actions of N and DL promoting EPC specification in all proneural cells. Sca might boost N and DL processing in the incipient EPCs while suppressing them or not affecting them in the incipient NPCs. Thus, it is possible that Sca or Sca-like molecule have a role in breaking the symmetrical actions of N and DL during certain lateral inhibition instances. It is also possible that Sca mediates long range N signaling during differentiation of some other tissues, either alone or in association with DL, as proposed by Renaud and Simpson [13]. By extending our results, it

might be possible to develop strong hypotheses for testing *in vivo*, cleanly sort the different activities of N, DL, and Sca, and understand the fascinating *in vivo* developmental mechanisms involving N, DL, and Sca.

Conclusion

Sca binds DL and suppresses a DL activity that is independent of DL's activity as a ligand of N. This DL activity requires the full length DL and is not enhanced by expression of just the DL intracellular domain, which is different from the mechanism underlying Notch activity. *Da* protein, *fng* RNA, and *pan* RNA responds positively to the N independent DL activity we have discovered. These could be direct or indirect targets. Our microarray analysis has identified many more putative targets of N independent DL activity that can be explored for a better understanding of the complex interactions among DL, Sca, and N during *Drosophila* development.

Methods

DNA constructs

Sca-Gfp: The stop codon of *sca* was replaced with a glycine codon and fused in-frame with GFP to obtain Sca-GFP. A Bam HI-KpnI fragment containing this *sca* sequence was cloned into pEGFP vector (Clontech). The XbaI fragment containing Sca-GFP coding fragment was cloned into the pCaSpeR-hs vector. DLΔI: A stop codon and a XbaI restriction site was introduced after the transmembrane domain using PCR. The PCR product was checked for mutations and used to replace the BstEII-BcgI fragment in the DL cDNA. An Eco RI-XbaI fragment from this construct (DL amino acid 1 to 620) was cloned into the pCaSpeR-hs vector. DLIC: The DL intracellular region (codon 619 to the stop codon 881) was PCR amplified, checked for errors, and cloned into the BglII-XbaI sites in the pCaSpeR-hs vector.

Cell lines and conditioned medium

S2-N, S2-DL, and S2-Sca cells have been previously described [14,20,49]. Other cell lines were established using the standard calcium phosphate transfection procedure and hygromycin selection. Conditioned medium was produced as described in Powell et al. [14], using serum-free or serum-containing Shields and Sang's M3 medium. For experiments, cells were heat shocked at 37°C for 30 minutes in a water bath, allowed to synthesize proteins for 2 hours, washed in culture medium without serum, mixed with the appropriate cell lines, and shaken gently in 14 ml falcon tubes for two hours or the indicated time. See Wesley and Mok [20] for more details.

Immunoprecipitations, western blotting, northern blotting, RNA in situ, and protein staining

Procedures described in Lieber et al. [43], Wesley [24], Wesley and Saez [18], and Wesley and Mok [20] were fol-

lowed. Eight per cent SDS-PAGE systems were used for western blotting, unless otherwise indicated; 1% formaldehyde-MOPS agarose system for northern blottings. *fringe* cDNA (from Dr. Ken Irvine), *rp49* cDNA, and rt-PCR amplified *pangolin* cDNA were used to prepare probes for northern blots. Incubation times with ligands were two hours for western blots and 45 minutes for all northern blots; it was three hours for *fng* and *pan* northern blot showing the effect of Sca (Fig. 4D).

Antibodies: α Sca (mAb sca1) and α DIEC (C594.9B) were obtained from the Developmental Studies Hybridoma Bank; α GFP (G-6539) and α Hsp70 (H-5147) from Sigma; α DIIC (GPC2) from Dr. Marc Muskavitch, α DIIC (dC-19) from Santa Cruz Biotechnology, α Da (DAM 109-10) from Dr. Claire Cronmiller; and α NI from Dr. Toby Lieber.

Microarray analysis

Heat shocked S2 and S2-DI cells were treated with Sca or non-Sca medium for 45 minutes before extracting RNA. GeneChip Drosophila Genome Arrays from Affymetrix were used. RNAs were extracted, checked, and processed for hybridization according to procedures suggested by Affymetrix. We pooled RNA from two independent experiments and used three such pooled samples as replicates for each treatment. The MicroArray Core Facility at the University of Vermont prepared the probes, hybridized the chips, and statistically analyzed the data (using the GeneSifter program). We used the Microarray Suite program to examine the data.

Authors' contributions

LPM and TQ designed and carried out many of the experiments in cultured cells and flies; BB carried out the immuno-precipitation experiments with cultured cells and helped in interpretation of data; MLC made the Sca GFP construct and assisted in many experiments; AH performed some experiments in cultured cells and prepared and maintained cell lines; FA assisted in statistical analyses and interpretation of data; and CSW conceived the study, designed experiments, and performed or participated in many of experiments in cultured cells and flies. LPM, BB, and MLC helped CSW in drafting the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Drs. Claire Cronmiller, Nick Baker, Toby Lieber, and Matt Rand for materials; Dr. Matt Rand and Uma Wesley for critical comments on the manuscript. We also thank the two reviewers and a member of the Editorial Board for helpful comments and advice. The work was supported by the grant R 01 NS43122-03 to C. S. Wesley.

References

1. Artavanis-Tsakonas S, Rand MD, Lake RJ: **Notch signaling: Cell fate control and signal integration in development.** *Science* 1999, **284**:770-776.

2. Mumm JS, Kopan R: **Notch Signaling: From the Outside.** *Dev Biol* 2000, **228**:151-165.
3. Cabrera CV: **Lateral inhibition and cell fate during neurogenesis in Drosophila: the interactions between scute, Notch and Delta.** *Development* 1990, **109**:733-742.
4. Jarman AP, Grell EH, Ackerman L, Jan LY, Jan YN: **Atonal is the proneural gene for Drosophila photoreceptors.** *Nature* 1994, **369**:398-400.
5. Vaessin H, Brand M, Jan LY, Jan YN: **daughterless is essential for neuronal precursor differentiation but not for initiation of neuronal precursor formation in Drosophila embryo.** *Development* 1994, **120**:935-945.
6. Brown NL, Paddock SW, Sattler CA, Cronmiller C, Thomas BJ, Carroll SB: **daughterless is required for Drosophila photoreceptor cell determination, eye morphogenesis, and cell cycle progression.** *Dev Biol* 1996, **179**:65-78.
7. Giebel B, Stuttem I, Hinz U, Campos-Ortega JA: **Lethal of Scute requires overexpression of daughterless to elicit ectopic neuronal development during embryogenesis in Drosophila.** *Mech Dev* 1997, **63**:75-87.
8. Cronmiller C and Cummings CC: **The daughterless gene product in Drosophila is a nuclear protein that is broadly expressed throughout the organism during development.** *Mech Dev* 1993, **42**:159-169.
9. Karp X, Greenwald I: **Post-transcriptional regulation of the E/Daughterless ortholog HLH-2, negative feedback, and birth order bias during the AC/VU decision in C. elegans.** *Genes Dev* 2003, **17**:3100-3111.
10. Cummings CA, Cronmiller C: **The daughterless gene functions together with Notch and Delta in the control of ovarian follicle development in Drosophila.** *Development* 1994, **120**:381-394.
11. Heitzler P, Simpson P: **The choice of cell fate in the epidermis of Drosophila.** *Cell* 1991, **64**:1083-1092.
12. Baker NE, Mlodzik M, Rubin GM: **Spacing differentiation in the developing Drosophila eye: A fibrinogen-related lateral inhibitor encoded by scabrous.** *Science* 1990, **250**:1370-1377.
13. Renaud O, Simpson P: **scabrous modifies epithelial cell adhesion and extends the range of lateral inhibition signaling during development of the spaced bristle pattern in Drosophila.** *Dev Biol* 2001, **240**:361-376.
14. Powell PA, Wesley C, Spencer S, Cagan RL: **Scabrous complexes with Notch to mediate boundary formation.** *Nature* 2001, **409**:626-630.
15. Lee E-C, Yu S-Y, Baker NE: **The Scabrous protein can act as an extracellular antagonist of Notch signaling in the Drosophila wing.** *Curr Biol* 2000, **10**:931-934.
16. Fehon RG, Kooh PJ, Rebay I, Regan CL, Xu T, Muskavitch M, Artavanis-Tsakonas S: **Molecular interaction between the protein products of the neurogenic loci Notch and Delta, two EGF-homologous genes in Drosophila.** *Cell* 1990, **61**:523-534.
17. Klueg KM, Muskavitch MAT: **Ligand-receptor interactions and trans-endocytosis of Delta, Serrate and Notch: members of the Notch signaling pathway in Drosophila.** *Development* 1999, **112**:3289-3297.
18. Wesley CS, Saez L: **Analysis of Notch lacking the carboxyl terminus identified in Drosophila embryos.** *J Cell Biol* 2000, **149**:683-696.
19. Mishra-Gorur K, Rand MD, Perez-Villamil B, Artavanis-Tsakonas S: **Down-regulation of Delta by proteolytic processing.** *J Cell Biol* 2002, **159**:313-324.
20. Wesley CS, Mok L-P: **Regulation of Notch signaling by a novel mechanism involving Suppressor of Hairless stability and carboxyl terminus-truncated Notch.** *Mol Cell Biol* 2003, **23**:5581-5593.
21. Ahimou F, Mok L-P, Bardot B, Wesley CS: **The adhesion force of Notch with Delta and the rate of Notch signaling.** *J Cell Bio* 2004, **167**:1217-1229.
22. Bardot B, Mok L-P, Thayer T, Ahimou F, Wesley CS: **The Notch amino terminus regulates protein levels and Delta-induced clustering of Drosophila receptors.** *Exp Cell Res* 2005, **304**:202-223.
23. Lee E-C, Baker NE: **GP300Sca is not a high affinity Notch ligand.** *Biochem. Biophysical Res Comm* 1996, **225**:720-725.
24. Wesley CS: **Notch and Wingless regulate expression of cuticle patterning genes.** *Mol Cell Biol* 1999, **19**:5743-5758.

25. Parks AL, Kleug KM, Stout JR, Muskavitch MA: **Ligand endocytosis drives receptor dissociation and activation in the Notch pathway.** *Development* 2000, **127**:1373-1385.
26. Struhl G, Adachi A: **Requirements for presenilin-dependent cleavage of notch and other transmembrane proteins.** *Mol Cell* 2000, **6**:625-636.
27. Pavlopoulos E, Pitsouli C, Kleug KM, Muskavitch MA, Moschonas NK, Delidakis C: **neuralized encodes a peripheral membrane protein involved in delta signaling and endocytosis.** *Dev Cell* 2001, **1**:807-816.
28. Overstreet E, Chen X, Wendland B, Fischer JA: **Either part of a Drosophila epsin protein, divided after the ENTH domain, functions in endocytosis of delta in the developing eye.** *Curr Biol* 2003, **13**:854-860.
29. Wang W, Struhl G: **Drosophila Epsin mediates a select endocytic pathway that DSL ligands must enter to activate Notch.** *Development* 2004, **131**:5367-5380.
30. Ikeuchi T, Sisodia S: **The Notch ligands, Delta 1 and Jagged 2, are substrates for Presenilin-dependent "γ-Secretase" cleavage.** *J Biol Chem* 2003, **278**:7751-7754.
31. Bland CE, Kimberly P, Rand MD: **Notch-induced proteolysis and Nuclear localization of the Delta ligand.** *J Biol Chem* 2003, **278**:13607-13610.
32. La Voie MJ, Selkoe DJ: **The Notch ligands, Jagged and Delta, are sequentially processed by alpha-Secretase and Presenilin/gamma-Secretase and release signaling fragments.** *J Biol Chem* 2003, **278**:34427-34437.
33. Six E, Ndiaye D, Laabi Y, Brou C, Gupta-Rossi N, Israel A, Logeat F: **The Notch ligand Delta 1 is sequentially cleaved by an ADAM protease and γ-secretase.** *Proc Nat Acad Sci USA* 2003, **100**:7638-7643.
34. Smith JEll, Cronmiller C: **The Drosophila daughterless gene autoregulates and is controlled by both positive and negative cis regulation.** *Development* 2001, **128**:4705-4714.
35. Irvine KD, Wieschaus E: **fringe, a boundary-specific signaling molecule, mediates interactions between dorsal and ventral cells during Drosophila wing development.** *Cell* 1994, **79**:595-606.
36. Moloney DJ, Panin VM, Johnston SH, Chen J, Shao L, Wilson R, Wang Y, Stanley P, Irvine KD, Haltiwanger RS, Vogt TF: **Fringe is a glycosyltransferase that modifies Notch.** *Nature* 2000, **406**:369-375.
37. Bruckner K, Perez L, Claussen H, Cohen S: **Glycosyltransferase activity of Fringe modulates Notch-Delta interactions.** *Nature* 2000, **406**:411-415.
38. Panin VM, Shao L, Lei L, Moloney DJ, Irvine KD, Haltiwanger RS: **Notch ligands are substrates for protein O-Fucosyltransferase-I and Fringe.** *J Biol Chem* 2002, **277**:29945-29952.
39. Brunner E, Peter O, Schweitzer L, Basler K: **pangolin encodes a Lef-1 homologue that acts downstream of Armadillo to transduce the Wingless signal in Drosophila.** *Nature* 1997, **385**:829-833.
40. van de Wetering M, Cavallo R, Dooijes D, van Beest M, van Es J, Loureiro J, Ypma Y, Hursh D, Jones T, Beijsovec A, Peifer M, Mortin M, Clevers H: **Armadillo coactivates transcription driven by the product of the Drosophila segment polarity gene dTCF.** *Cell* 1997, **88**:789-799.
41. Couso JP, Martinez-Arias A: **Notch is required for wingless signaling in the epidermis of Drosophila.** *Cell* 1994, **67**:311-323.
42. Rulifson EJ, Blair SS: **Notch regulates wingless expression and is not required for reception of the paracrine wingless signal during wing margin neurogenesis in Drosophila.** *Development* 1995, **121**:2813-2824.
43. Cadigan KM, Nusse R: **wingless signaling in the Drosophila eye and embryonic epidermis.** *Development* 1996, **122**:2801-2812.
44. Axelrod JD, Matsuno K, Artavanis-Tsakonas S, Perrimon N: **Interaction between Wingless and Notch signaling pathways mediated by Dishevelled.** *Science* 1996, **271**:1826-1832.
45. Fortini ME, Rebay I, Caron LA, Artavanis-Tsakonas S: **An activated Notch receptor blocks cell-fate commitment in the developing Drosophila eye.** *Nature* 1993, **365**:555-557.
46. Struhl G, Fitzgerald K, Greenwald I: **Intrinsic activity of the Lin-12 and Notch intracellular domains in vivo.** *Cell* 1993, **74**:331-345.
47. Lieber T, Kidd S, Alcamo E, Corbin V, Young MW: **Antineurogenic phenotypes induced by truncated Notch proteins indicate a role in signal transduction and may point to a novel function for Notch in nuclei.** *Genes Dev* 1993, **7**:1949-1965.
48. Rebay I, Fleming RJ, Fehon RG, Cherbas L, Cherbas P, Artavanis-Tsakonas S: **Specific EGF repeats of Notch mediate interactions with Delta and Serrate: implications for Notch as a multi-functional receptor.** *Cell* 1991, **67**:687-699.
49. Lieber T, Wesley CS, Alcamo E, Hassel B, Krane JF, Campos-Ortega JA, Young MW: **Single amino acid substitutions in EGF-like elements of Notch and Delta modify Drosophila development and affect cell adhesion in vitro.** *Neuron* 1992, **9**:847-859.
50. Huppert SS, Jacobsen TL, Muskavitch MAT: **Feedback regulation is central to Delta-Notch signaling required for Drosophila wing vein morphogenesis.** *Development* 1997, **124**:3283-3291.
51. de Celis JF, Bray S, Garcia-Bellido A: **Notch signaling regulates veinlet expression and establishes boundaries between veins and interveins in the Drosophila wing.** *Development* 1997, **124**:1919-1928.
52. Ascano JM, Beverly LJ, Capobianco AJ: **The C-terminal PDZ-ligand of JAGGED 1 is essential for cellular transformation.** *J Biol Chem* 2003, **278**:8771-8779.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

